

Supplementary Table 3. Baseline characteristics, time to evaluation, and time to thrombolysis according to the ambulance prenotification

	Prenotification (-) (n=34)	Prenotification (+) (n=13)	<i>P</i>
Age (years)	70.0 (62.8–74.3)	68.0 (64.5–74.0)	0.802
Sex, male			0.324
Initial NIHSS	8.0 (3.8–16.3)	10.0 (8.5–13.5)	0.535
Stroke onset-to-door (min)	97.0 (45.3–152.8)	89.0 (29.0–218.0)	0.887
Door-to-group messaging (min)	7.0 (4.0–12.0)	3.0 (2.0–11.5)	0.028
Door-to-CT (min)	17.0 (12.0–20.0)	7.0 (5.0–9.0)	<0.001
Door-to-MRI (min)	38.0 (30.0–62.0)	34.0 (30.5–46.5)	0.403
Door-to-CBC (min)	49.0 (30.5–61.5)	32.0 (22.0–43.0)	0.016
Door-to-PT (routine) (min)	46.0 (39.0–63.5)	34.0 (30.5–47.5)	0.018
Door-to-PT (POCT) (min)	10.0 (7.0–20.5)	2.0 (1.0–4.0)	<0.001
Thrombolysis type			
IVT only	22 (64.7)	6 (46.2)	0.324
IAT only	11 (32.4)	2 (15.4)	0.301
IVT plus IAT	1 (2.9)	5 (38.5)	0.004
Door-to-IVT (min)	26.0 (20.0–36.0)	15.0 (12.0–19.0)	<0.001
Door-to-IAT (min)	179.0 (46.0–137.5)	89.5 (78.3–99.3)	0.473
Stroke onset-to-IVT (min)	113.0 (80.0–152.0)	57.0 (45.0–158.0)	0.363
Stroke onset-to-IAT (min)	227.5 (174.8–281.5)	252.0 (112.0–655.3)	1.000
CT-to-IVT (min)	10.5 (6.0–16.5)	6.0 (6.0–9.0)	0.157
IVT-to-IAT (min)	78.0 (78.0–78.0)	67.5 (63.5–101.3)	0.857
CT-to-MRI (min)	21.0 (12.5–39.8)	28.0 (24.5–41.0)	0.198
CT-to-IAT (min)	76.5 (35.0–143.5)	82.0 (71.0–93.8)	0.798
MRI-to-IAT (min)	60.0 (49.5–74.8)	61.5 (38.0–72.3)	0.959

NIHSS, the NIH Stroke Scale; CT, computed tomography; MRI, magnetic resonance imaging; CBC, complete blood count; PT, prothrombin time; POCT, point of care test; IVT, intravenous thrombolysis; IAT, intra-arterial thrombectomy.