

**Supplementary Table 2.** Absolute (minutes) and relative (%) time spent with SBP and MAP >120% or <80% of the reference value, overall and by mRS score at 3 months

Variable	All patients	mRS 0–2	mRS 3–6	Missing mRS
Number	164	75	80	9
SBP >120% of reference (min)	8.6 (0–158)	8.6 (0–84)	8.6 (0–158)	7.3 (0–59)
MAP >120% of reference (min)	9.5 (0–223)	9.2 (0–127)	9.7 (0–223)	10.7 (0–63)
SBP <80% of reference (min)	8.9 (0–152)	9.0 (0–143)	8.9 (0–152)	7.6 (0–9)
MAP <80% of reference (min)	7.2 (0–160)	7.8 (0–75)	7.3 (0–160)	1 (0–1)
SBP >120% of reference (%)	10.9 (0–93)	12.1 (0–93)	9.3 (0–84)	16.3 (0–82)
MAP >120% of reference (%)	11.9 (0–97)	13.1 (0–89)	10.3 (0–97)	19.5 (0–88)
SBP <80% of reference (%)	9.6 (0–92)	12.3 (0–90)	8.2 (0–92)	6.6 (0–9.6)
MAP <80% of reference (%)	7.8 (0–89)	10.6 (0–89)	6.6 (0–82)	1 (0–1)

Values are presented as geometric mean (range).

SBP, systolic blood pressure; MAP, mean arterial pressure; mRS, modified Rankin Scale.