

**Supplementary Table 3.** Comparisons in clinical outcomes according to use of intravenous thrombolysis prior to endovascular treatment using the stabilized inverse propensity score of MT+IVT as weight into logistic regression model

Outcomes	OR (95% CI)	P
Clinical outcomes		
Favorable outcome	1.37 (1.09–1.73)	0.008
Excellent outcome	1.71 (1.33–2.21)	<0.001
90-Day mortality	0.78 (0.60–1.02)	0.065
Hemorrhagic complications		
Any ICH	1.07 (0.85–1.36)	0.550
sICH	1.19 (0.77–1.82)	0.430
Procedural outcomes		
Successful reperfusion	1.56 (1.22–1.99)	<0.001
Excellent reperfusion	1.09 (0.87–1.36)	0.440
Rescue therapy	0.80 (0.61–1.04)	0.100
Emboli uninvolved territory	0.69 (0.45–1.04)	0.080
Vessel perfusion	1.05 (0.64–1.73)	0.850
Vasospasme	0.57 (0.25–1.32)	0.190
Number of passes (>2)	0.79 (0.64–0.99)	0.037
Groin puncture to successful reperfusion time (45 min)	1.31 (1.05–1.63)	0.017

IMT, mechanical thrombectomy; IVT, intravenous thrombolysis; OR, odds ratio; CI, confidence interval; ICH, intracerebral hemorrhage; sICH, symptomatic intracerebral hemorrhage.