

Supplementary Table 13. Net clinical benefit of the combination of ischemic stroke and systemic embolism and ICH

Type of diagnosis	TE					ICH				
	Not receiving OACs		Receiving exposure to all OACs		Difference in rate of TE (off-on)	Not receiving OACs		Receiving exposure to all OACs		Difference in rate of ICH (on-off)
	PY	Annual rate per 100 PY	PY	Annual rate per 100 PY		PY	Annual rate per 100 PY	PY	Annual rate per 100 PY	
Total cohort	331.9	3.3	235.0	2.6	0.7	315.7	11.4	229.5	5.2	-6.2
Non-traumatic diagnosis	193.4	3.6	113.0	0.9	2.7	185.7	12.4	109.6	3.6	-8.8
Traumatic diagnosis	138.4	2.9	122.0	4.1	-1.5	130.0	10.0	119.9	6.7	-3.3
Annual net clinical benefit	Weight=1.5				Weight=1.0	Weight=2.0				
Total cohort	10.3 (10.1–10.5)				7.1 (6.9–7.3)	13.5 (13.2–13.8)				
Non-traumatic diagnosis	16.3 (16.0–16.6)				11.8 (11.5–12.0)	20.8 (20.4–21.2)				
Traumatic diagnosis	3.8 (3.4–4.2)				2.1 (1.9–2.4)	5.5 (5.0–6.0)				

Values are presented as annual net clinical benefit (95% confidence interval).

ICH, intracranial hemorrhage; TE, thromboembolism; OAC, oral anticoagulant; PY, person-year.