

**Supplementary Table 6.** Adjusted hazard ratios of clinical outcomes with a quarantine period of 6 weeks with binary time-dependent exposure (full cohort)

	Stroke/SE/death (185 events)	Stroke/SE (17 events)	Death (174 events)	Recurrent intracranial bleeding (48 events)	Major extracranial bleeding (21 events)
Exposure to OAC					
No treatment (reference)					
Treatment	0.10 (0.05–0.21)	0.56 (0.16–1.96)	0.07 (0.03–0.17)	0.43 (0.19–0.97)	1.73 (0.71–4.20)
Age (yr)					
65–74 (reference)					
≥75	2.21 (1.13–4.33)		2.64 (1.25–5.54)	0.45 (0.21–0.94)	2.41 (0.32–18.10)
Sex					
Male (reference)					
Female	0.69 (0.49–0.95)	6.24 (1.43–27.28)	0.61 (0.44–0.86)	1.20 (0.67–2.16)	2.28 (0.88–5.90)
Prior stroke/TIA/SE*					
Prior stroke/TIA/SE*	1.21 (0.68–2.14)	0.73 (0.10–5.49)	1.24 (0.70–2.21)		
Prior major bleeding (except ICH) *					
Prior major bleeding (except ICH) *	0.94 (0.57–1.55)		0.89 (0.53–1.50)	1.98 (0.81–4.83)	4.27 (1.65–11.05)
Chronic heart failure*					
Chronic heart failure*	1.40 (1.01–1.94)		1.53 (1.09–2.15)		
Peripheral artery disease*					
Peripheral artery disease*	0.98 (0.69–1.39)		1.04 (0.73–1.50)		
Chronic kidney disease*					
Chronic kidney disease*	1.30 (0.92–1.85)		1.26 (0.88–1.80)		
Hypertension*					
Hypertension*	0.90 (0.57–1.40)		0.85 (0.54–1.35)	0.82 (0.37–1.81)	
Dyslipidemia*					
Dyslipidemia*	0.94 (0.66–1.35)		0.91 (0.63–1.32)		
Diabetes*					
Diabetes*	1.23 (0.89–1.71)		1.24 (0.88–1.75)	0.63 (0.30–1.31)	
Beta-blockers*					
Beta-blockers*	1.18 (0.87–1.61)		1.25 (0.90–1.72)		
Statins*					
Statins*	0.73 (0.51–1.04)		0.69 (0.48–0.99)		
Proton-pump inhibitors*					
Proton-pump inhibitors*	1.44 (1.05–1.97)		1.47 (1.06–2.03)		
Digoxin*					
Digoxin*	1.53 (1.11–2.11)		1.47 (1.05–2.06)		
Score Charlson					
<4 (reference)					
≥4	1.33 (0.89–1.99)		1.27 (0.84–1.93)	1.06 (0.57–1.96)	
CHA <sub>2</sub> DS <sub>2</sub> -VASc score					
<3 (reference)					
≥3	0.93 (0.57–1.52)		0.89 (0.53–1.47)		
HAS-BLED score					
<3 (reference)					
≥3	1.03 (0.72–1.49)		1.12 (0.77–1.64)	0.87 (0.45–1.68)	

Values are presented as hazard ratio (95% confidence interval).

SE, systemic embolism; OAC, oral anticoagulant; TIA, transient ischemic attack; ICH, intracranial hemorrhage.

\*1=yes vs. 0=no.