

**Supplementary Table 7.** Adjusted hazard ratios of clinical outcomes with a quarantine period of 6 weeks with binary time-dependent exposure and categorical exposure

	Stroke/SE/death (185 events)	Stroke/SE (17 events)	Death (174 events)	Recurrent intracranial bleeding (48 events)	Major extracranial bleeding (21 events)
<b>Exposure to OACs</b>					
0% (reference)					
% <90+current not exposed	2.43 (1.56–3.78)	2.55 (0.57–11.45)	2.43 (1.55–3.82)	1.00 (0.30–3.26)	3.36 (0.92–12.22)
% <90+current exposed	0.25 (0.10–0.60)	-	0.26 (0.11–0.64)	0.50 (0.15–1.62)	2.41 (0.76–7.60)
% ≥90	0.10 (0.04–0.24)	0.87 (0.24–3.11)	0.04 (0.01–0.17)	0.38 (0.13–1.08)	1.01 (0.28–3.68)
<b>Age (yr)</b>					
65–74 (reference)					
≥75	2.38 (1.21–4.67)		2.85 (1.35–6.01)	0.47 (0.23–0.99)	
<b>Sex</b>					
Male (reference)					
Female	0.67 (0.48–0.94)	6.29 (1.44–27.50)	0.60 (0.43–0.85)	1.20 (0.67–2.14)	
<b>Prior stroke/TIA/SE*</b>					
Prior stroke/TIA/SE*	1.18 (0.67–2.09)		1.21 (0.68–2.15)		
<b>Prior major bleeding (except ICH)*</b>					
Prior major bleeding (except ICH)*	0.88 (0.53–1.46)		0.84 (0.50–1.41)	2.04 (0.83–5.01)	3.70 (1.42–9.63)
<b>Chronic heart failure*</b>					
Chronic heart failure*	1.39 (1.00–1.92)		1.52 (1.09–2.14)		
<b>Peripheral artery disease*</b>					
Peripheral artery disease*	0.96 (0.68–1.37)		1.03 (0.72–1.48)		
<b>Chronic kidney disease*</b>					
Chronic kidney disease*	1.33 (0.94–1.89)		1.29 (0.89–1.85)		
<b>Hypertension*</b>					
Hypertension*	0.90 (0.58–1.41)		0.85 (0.54–1.35)		
<b>Dyslipidemia*</b>					
Dyslipidemia*	0.89 (0.62–1.27)		0.85 (0.59–1.24)		
<b>Diabetes*</b>					
Diabetes*	1.27 (0.91–1.76)		1.28 (0.91–1.80)		
<b>Beta-blockers*</b>					
Beta-blockers*	1.17 (0.86–1.60)		1.24 (0.89–1.71)		
<b>Statins*</b>					
Statins*	0.72 (0.50–1.03)		0.68 (0.47–0.98)		
<b>Proton-pump inhibitors*</b>					
Proton-pump inhibitors*	1.40 (1.02–1.92)		1.42 (1.02–1.97)		
<b>Digoxin*</b>					
Digoxin*	1.49 (1.08–2.05)		1.43 (1.02–2.01)		
<b>Score Charlson</b>					
<4 (reference)					
≥4	1.32 (0.88–1.97)		1.26 (0.83–1.92)	0.90 (0.51–1.61)	
<b>CHA<sub>2</sub>DS<sub>2</sub>-VASC score</b>					
<3 (reference)					
≥3	0.93 (0.57–1.53)		0.89 (0.53–1.47)		
<b>HAS-BLED</b>					
<3 (reference)					
≥3	1.05 (0.73–1.52)		1.14 (0.78–1.66)	0.82 (0.43–1.57)	

Values are presented as hazard ratio (95% confidence interval).

SE, systemic embolism; OAC, oral anticoagulant; TIA, transient ischemic attack; ICH, intracranial hemorrhage.

\*1=yes vs. 0=no.