

Supplementary Table 8. Adjusted hazard ratios of all-cause mortality with a quarantine period of 6 weeks with binary time-dependent exposure and time-dependent covariates

	Death (174 events)
Exposure to OACs	
No exposure (reference)	
Exposure	0.07 (0.03–0.17)
Stroke/SE*	3.75 (1.58–8.91)
Recurrent intracranial bleeding*	1.95 (1.13–3.37)
Major extracranial bleeding*	1.96 (0.89–4.29)
Age (yr)	
65–74 (reference)	
≥75	2.67 (1.26–5.62)
Sex	
Male (reference)	
Female	0.57 (0.41–0.81)
Prior stroke/TIA/SE*	1.20 (0.67–2.15)
Prior major bleeding (except ICH)*	0.84 (0.50–1.41)
Chronic heart failure*	1.61 (1.15–2.26)
Peripheral artery disease*	1.06 (0.74–1.53)
Chronic kidney disease*	1.23 (0.85–1.77)
Hypertension*	0.81 (0.51–1.29)
Dyslipidemia*	0.90 (0.62–1.31)
Diabetes*	1.22 (0.87–1.72)
Beta-blockers*	1.22 (0.88–1.70)
Statins*	0.68 (0.47–0.98)
Proton-pump inhibitors*	1.45 (1.05–2.01)
Digoxin*	1.43 (1.02–2.00)
Score Charlson	
<4 (reference)	
≥4	1.24 (0.81–1.89)
CHA ₂ DS ₂ -VASc score	
<3 (reference)	
≥3	0.91 (0.55–1.51)
HAS-BLED	
<3 (reference)	
≥3	1.17 (0.80–1.71)

Values are presented as hazard ratio (95% confidence interval).

OAC, oral anticoagulant; SE, systemic embolism; TIA, transient ischemic attack; ICH, intracranial hemorrhage.

*1=yes vs. 0=no.