Study or subgroup	Recanalization Mean	SD	Total	No recanalization Mean	SD	Total	Weight	SMD IV, random, 95% CI
Delgado-Mederos et al., 2008	146.9	24.3	44	151.1	31.9	36	20.4%	-0.15 [-0.59, 0.29]
Molina et al., 2009	150	27	17	157	21	18	8.9%	-0.28 [-0.95, 0.38]
Tsivgoulis et al., 2007	152	27	94	157	21	257	70.7%	-0.22 [-0.46, 0.02]
Total (95% CI)			**155**			**311**	**100.0%**	**-0.21 [-0.41, -0.01]**

Heterogeneity: Tau^2=0.00; Chi^2=0.13, df=2 (*P*=0.94); I^2=0%
Test for overall effect: Z=2.07 (*P*=0.04)



Study or subgroup	Recanalization Mean	SD	Total	No recanalization Mean	SD	Total	Weight	SMD IV, random, 95% CI
Delgado-Mederos et al., 2008	79.5	12.2	44	84.8	19.3	36	69.2%	-0.33 [-0.78, 0.11]
Molina et al., 2009	76	14	17	80	18	18	30.8%	-0.24 [-0.91, 0.42]
Total (95% CI)			**61**			**54**	**100.0%**	**-0.30 [-0.67, 0.06]**

Heterogeneity: Tau^2=0.00; Chi^2=0.05, df=1 (*P*=0.82); I^2=0%
Test for overall effect: Z=1.62 (*P*=0.11)



Supplementary Figure 12. Forest plots evaluating the association of pre-treatment (A) systolic blood pressure levels and (B) diastolic blood pressure levels with recanalization. SMD, standardized mean difference; SD, standard deviation; IV, intravenous; CI, confidence interval.