

Supplementary Table 1. Univariate analysis of outcome predictors (mRS 0–3)

Variable	Unfavorable (n=99)	Favorable (n=73)	P
Age (yr)	72.9±14.2	63.8±15.5	<0.001
Female sex	36 (36.4)	26 (35.6)	1
Dyslipidemia	49 (49.5)	20 (27.4)	0.003
Diabetes mellitus	20 (20.2)	6 (8.2)	0.005
Tobacco use (current or past)	28 (28.3)	28 (38.4)	0.187
Hypertension	65 (65.7)	32 (43.8)	0.005
NIHSS	19.3±4.5	17.5±4.3	0.011
Left sided stroke	46 (46.5)	29 (39.7)	0.355
iv tPA	56 (56.6)	49 (67.1)	0.205
Drip and ship	22 (22.2)	11 (15.1)	0.327
ICA occlusion	21 (21.2)	11 (15.1)	0.329
Delay till imaging (min)	167.3±137.9	153.1±116.4	0.468
Volume T max <6 sec (mL)	185.1±68	154.7±68.4	0.004
Mismatch ratio	1.8±0.7	1.7±0.6	0.166
Core volume (mL)	108.4±41.7	94±26.6	<0.001
Received MT	74 (74.7)	56 (76.7)	0.851

Values are presented as mean±standard deviation or number (%).

mRS, modified Rankin Scale; NIHSS, National Institute of Health Stroke Scale; iv tPA, intravenous tissue plasminogen activator; ICA, internal carotid artery; MT, mechanical thrombectomy.

Supplementary Table 2. Univariable analysis of 90-day mortality

Variable	Dead (n= 54)	Alive (n=118)	P
Age (yr)	77.63±11.99	65.09±15.21	<0.001
Female sex	19 (35.2)	43 (36.4)	1
Dyslipidemia	27 (50)	42 (35.6)	0.094
Diabetes	15 (27.8)	11 (9.3)	0.003
Hypertension	37 (68.5)	60 (50.8)	0.033
NIHSS	19.78±4.7	17.96±4.25	0.019
Tobacco use	11 (20.4)	45 (38.1)	0.023
ICA occlusion	10 (18.5)	22 (18.6)	1.000
Delay till imaging (min)	164.87±155.96	159.61±115.33	0.825
Drip and ship	12 (22.2)	21 (17.8)	0.534
iv tPA	30 (55.6)	75 (63.6)	0.401
Volume core (mL)	111.3±43.4	98.11±32.49	0.025
Volume T max >6 sec (min)	192.02±64.11	163.13±70.38	0.009
Mismatch ratio	1.86±0.75	1.7±0.65	0.173
MT	41 (31.5)	89 (68.5)	vs. control, 1.000
Control	13 (30.9)	29 (69.1)	

Values are presented as number (%) or mean±standard deviation.

NIHSS, National Institute of Health Stroke Scale; ICA, internal carotid artery; iv tPA, intravenous tissue plasminogen activator; MT, mechanical thrombectomy.