

Supplementary Table 3. Multivariable model of 90-day mortality predictors (pre specified adjustment for core volume, mismatch ratio, and treatment group)

Variable	aOR	95% CI	P
Age	1.11	1.07–1.16	<0.001
Diabetes	4.34	1.54–12.27	0.006
Volume core (each 10 mL)	1.32	1.16–1.53	<0.001
Core perfusion mismatch	2.38	1.26–4.47	0.006
MT vs. Control	0.49	0.19–1.37	0.177

aOR, adjusted odds ratio; CI, confidence interval; MT, mechanical thrombectomy.

Supplementary Table 4. Univariable analysis of sICH (ECASS II) predictors

Variable	sICH (n=31)	No sICH (n=130)	P
Age (yr)	72.89±12.15	67.74±15.98	0.051
Female sex	9 (29.0)	50 (38.5)	0.409
Dyslipidemia	12 (38.7)	51 (39.2)	1.000
Diabetes mellitus	9 (29.0)	14 (10.8)	0.002
Hypertension	22 (71.0)	67 (51.5)	0.068
NIHSS	18.53±3.46	18.6±4.67	0.925
Tobacco use	6 (19.4)	45 (34.6)	0.133
ICA occlusion	7 (22.6)	22 (16.9)	0.444
Delay till imaging (min)	149.71±113.66	163.7±131.43	0.548
Drip and ship	7 (22.6)	24 (18.5)	0.609
iv tPA	19 (61.3)	78 (60.0)	1.000
Volume core (mL)	112.43±38.47	100.6±35.67	0.126
Volume T max >6 sec (mL)	188.75±59.87	171.58±72.5	0.175
Mismatch ratio	1.8±0.72	1.76±0.7	0.795
MT (P for vs. control)	26 (20.6)	100 (76.4)	0.475
Control	5 (14.3)	30 (85.7)	

Values are presented as number (%) or mean±standard deviation.

sICH, symptomatic intracerebral hemorrhage; ECASS, European Cooperative Acute Stroke Study; NIHSS, National Institute of Health Stroke Scale; ICA, internal carotid artery; iv tPA, intravenous tissue plasminogen activator; MT, mechanical thrombectomy.