

Supplementary Table 2. Logistic regression for good outcome in anterior circulation stroke patients at follow-up

Variable	OR	95% CI	P
Age	0.95	0.95–0.96	<0.001
Sex	0.99	0.84–1.15	0.859
pmRS	0.51	0.47–0.56	<0.001
NIHSS at admission	0.88	0.86–0.89	<0.001
Treatment with IVT	1.57	1.35–1.84	<0.001
Successful reperfusion (mTICI 2b/3)	5.41	4.15–7.05	<0.001
General anesthesia	0.86	0.74–0.99	0.041
ICH	0.25	0.20–0.31	<0.001
Time groin to flow restoration	0.99	0.99–1.00	0.009

OR, odds ratio; CI, confidence interval; pmRS, pre-stroke modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; IVT, intravenous thrombolysis; mTICI, modified thrombolysis in cerebral infarction; ICH, intracerebral hemorrhage.

Supplementary Table 3. Logistic regression for mortality in anterior circulation stroke patients at follow-up

Variable	OR	95% CI	P
Age	1.07	1.06–1.07	<0.001
Sex	0.88	0.75–1.04	0.126
pmRS	1.37	1.29–1.46	<0.001
NIHSS at admission	1.10	1.09–1.12	<0.001
Treatment with IVT	0.67	0.57–0.79	<0.001
Successful reperfusion (mTICI 2b/3)	0.28	0.23–0.34	<0.001
General anesthesia	1.38	1.19–1.61	<0.001
ICH	2.73	2.26–3.31	<0.001
Time admission to groin	1.00	1.00–1.00	0.323

OR, odds ratio; CI, confidence interval; pmRS, pre-stroke modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; IVT, intravenous thrombolysis; mTICI, modified thrombolysis in cerebral infarction; ICH, intracerebral hemorrhage.