

Supplementary Table 5. Adjusted mean concentration of lipids, lipoproteins, apolipoproteins, and their ratios in cases and controls

Variable	Ischemic stroke (n=9,169)	Intracerebral hemorrhage (n=2,729)	Controls (n=11,898)
Total cholesterol (mmol/L)	4.61 (4.55–4.67)	4.56 (4.47–4.64)	4.72 (4.65–4.78)
HDL-C (mmol/L)	1.12 (1.10–1.14)	1.27 (1.24–1.29)	1.20 (1.18–1.22)
Non-HDL-C (mmol/L)	3.49 (3.43–3.55)	3.29 (3.21–3.37)	3.52 (3.46–3.58)
ApoA1 (g/L)	1.27 (1.25–1.29)	1.33 (1.31–1.35)	1.38 (1.36–1.40)
ApoB (g/L)	0.99 (0.97–1.01)	0.95 (0.93–0.97)	0.97 (0.95–0.99)
Total cholesterol/HDL-C ratio	4.53 (4.43–4.63)	3.91 (3.78–4.04)	4.34 (4.23–4.44)
ApoB/apoA1 ratio	0.84 (0.82–0.86)	0.77 (0.75–0.79)	0.76 (0.74–0.78)
Non-HDL-C/HDL-C ratio	3.53 (3.43–3.63)	2.91 (2.78–3.04)	3.34 (3.23–3.44)

Values are presented as least square mean (95% confidence interval) after adjustment for age, sex, region, smoking, waist-to-hip ratio, diabetes, physical activity, alternate healthy eating index, alcohol intake, psychosocial factors, and cardiac risk factors.

HDL-C, high-density lipoprotein cholesterol; Apo, apolipoprotein.