Supplementary Table 5. Continued

	pmRS 3–5 (n=92)	pmRS ≤2 (n=513)	*P*
Type of anesthesia			0.147
GA	67/91 (73.6)	403/510 (79.0)	
CS	24/91 (26.4)	94/510 (18.4)	
Switch local to GA	0/91 (0.0)	13/510 (2.6)	
Recanalization attempts	2 (1–3)	2 (1–3)	0.954

Values are presented as mean±standard deviation, median (interquartile range), or n (%).
ASPECTS, Alberta Stroke Program Early CT Score; pmRS, premorbid modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; BP, blood pressure; CT, computed tomography; MRI, magnetic resonance imaging; ICA, internal carotid artery; GA, general anesthesia; CS, conscious sedation.
*Significant differences; †NIHSS at admission was assessed based on the neurological symptoms observed at the time of admission, regardless of any preexisting physical deficits from prior strokes; ‡Included medication with aspirin and/or clopidogrel; §Included medication with either apixaban, rivaroxaban, dabigatran, edoxaban, or phenprocoumon.



Supplementary Figure 1. Flowchart of patient selection. GSR-ET, German Stroke Registry–Endovascular Treatment; pmRS, premorbid modified Rankin Scale; BA, basilar artery; VA, vertebral artery; PCA, posterior cerebral artery; ACA, anterior cerebral artery; eCA, extracranial carotid artery; ET, endovascular treatment; ASPECTS, Alberta Stroke Program Early CT Score.