Supplementary Table 2. Premorbid ASPECTS ≤5 patients stratified by effective ET

	Effective ET (n=18, 19.6%)	Futile ET (n=74, 80.4%)	*P*
Baseline characteristics			
Age (yr)	75.9±13.4	81.6±11.3	0.069
Female sex	9/18 (50.0)	57/74 (77.0)	0.038*
pmRS	4 (3–4)	3 (3–4)	0.136
NIHSS	14 (12–21)	18 (15–21)	0.034*
BP systolic (mm Hg)	150 (137–170)	150 (129–169)	0.962
BP diastolic (mm Hg)	80 (71–91)	80 (70–90)	0.847
Heart rate (bpm)	86 (64–114)	85 (70–99)	0.927
Vascular risk factors			
Arterial hypertension	10/18 (55.6)	53/74 (71.6)	0.157
Diabetes	2/18 (11.1)	16/74 (21.6)	0.509
Dyslipidemia	6/18 (33.3)	22/73 (30.1)	0.782
Atrial fibrillation	6/18 (33.3)	40/72 (55.6)	0.117
Active smoking	1/14 (7.1)	4/68 (5.9)	0.858
Ship	4/18 (22.2)	24/74 (32.4)	0.570
Mothership	14/18 (77.8)	50/744 (67.2)	0.570
Workflow times (h)			
Last known normal to admission	3.6 (1.6–11.9)	3.6 (1.5–7.9)	0.757
Admission to groin puncture	1.3 (1.0–1.6)	1.3 (0.8–1.9)	0.859
Groin puncture to recanalization	0.6 (0.4–1.2)	0.6 (0.3–1.1)	0.402
Last known normal to recanalization	5.3 (3.3–17.4)	5.8 (3.8–10.6)	0.926
Stroke, imaging, and treatment characteristics			
ASPECTS	5 (4–5)	5 (3–5)	0.139
ASPECTS assessed on			
CT	15/18 (83.3)	71/74 (95.9)	0.251
MRI	3/18 (16.7)	3/74 (4.1)	0.251
Carotid-T	4/18 (22.2)	19/74 (25.7)	0.762
ICA (excluding carotid-T)	2/18 (11.1)	5/74 (6.8)	0.619
M1 proximal	8/18 (44.4)	34/74 (45.9)	0.909
M1 distal	3/18 (16.7)	11/74 (14.9)	0.849
M2	1/18 (5.6)	5/74 (6.8)	0.853
Left-sided occlusion	7/18 (38.9)	33/73 (45.2)	0.792
Intravenous thrombolysis	7/18 (38.9)	27/74 (36.5)	0.850
Type of anesthesia			
GA	15/18 (83.3)	52/73 (71.2)	0.381
CS	3/18 (16.7)	21/73 (28.8)	0.381
Switch local to GA	0/18 (0.0)	0/73 (0.0)	>0.999
Recanalization attempts	2 (1–4)	2 (1–3)	0.817
Clinical and radiological outcomes			
NIHSS at 24 h	14 (9–21)	20 (16–26)	0.002*
sICH	0/18 (0.0)	7/74 (9.5)	0.338
Procedure-related complications	2/18 (11.1)	13/74 (17.6)	0.727

Values are presented as mean±standard deviation, median (interquartile range), or n (%).
ET, endovascular treatment; ASPECTS, Alberta Stroke Program Early CT Score; pmRS, premorbid modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; BP, blood pressure; CT, computed tomography; MRI, magnetic resonance imaging; ICA, internal carotid artery; GA, general anesthesia; CS, conscious sedation; sICH, symptomatic intracranial hemorrhage.
*Significant differences.

https://doi.org/10.5853/jos.2025.00668