Supplementary Table 3. Premorbid ASPECTS ≤5 patients stratified by 90-day mortality

	Survival (n=27, 29.3%)	Death (n=65, 70.7%)	*P*
Baseline characteristics			
Age (yr)	77 (71–82)	83 (76–91)	0.011*
Female sex	14/27 (51.9)	52/65 (80.0)	0.010*
pmRS	3 (3–4)	3 (3–4)	0.501
NIHSS	16 (12–21)	18 (15–21)	0.043*
BP systolic (mm Hg)	150 (137–171)	150 (128–168)	0.564
BP diastolic (mm Hg)	80 (69–90)	80 (71–99)	0.260
Heart rate (bpm)	85 (71–98)	83 (66–120)	0.906
Vascular risk factors			
Arterial hypertension	17/27 (63.0)	46/63 (73.0)	0.452
Diabetes	3/27 (11.1)	15/64 (23.4)	0.252
Dyslipidemia	10/27 (37.0)	18/64 (28.1)	0.459
Atrial fibrillation	13/27 (48.1)	33/63 (52.4)	0.819
Active smoking	3/23 (13.0)	2/59 (3.4)	0.131
Ship	7/27 (25.9)	21/65 (32.3)	0.625
Mothership	20/27 (74.1)	44/65 (67.7)	0.625
Workflow times (h)			
Last known normal to admission	3.9 (2.3–10.3)	3.4 (1.4–7.4)	0.180
Admission to groin puncture	1.4 (1.1–1.8)	1.3 (0.8–1.8)	0.305
Groin puncture to recanalization	0.7 (0.4–1.2)	0.5 (0.3–1.0)	0.159
Last known normal to recanalization	7.2 (3.8–15.2)	5.7 (3.4–9.6)	0.156
Stroke, imaging, and treatment characteristics			
ASPECTS	5 (4–5)	5 (4–5)	0.207
ASPECTS assessed on			
CT	24/27 (88.9)	63/65 (96.9)	0.149
MRI	3/27 (11.1)	2/65 (3.1)	0.149
Carotid-T	5/27 (18.5)	18/65 (27.7)	0.435
ICA (excluding carotid-T)	3/27 (11.1)	4/65 (6.2)	0.414
M1 proximal	10/27 (37.0)	32/65 (49.2)	0.360
M1 distal	6/27 (22.2)	8/65 (12.3)	0.338
M2	3/27 (11.1)	3/65 (4.6)	0.354
Intravenous thrombolysis	7/27 (25.9)	27/65 (41.5)	0.235
Type of anesthesia			0.261
GA	20/26 (76.9)	47/65 (72.3)	
CS	6/26 (23.1)	18/65 (27.7)	
Switch local to GA	0 (0.0)	0 (0.0)	
Recanalization attempts (n)	2 (1–4)	2 (1–3)	0.366
Clinical and radiological outcomes			
NIHSS at 24 h	16 (11–21)	21 (17–26)	0.001*
sICH	0/27 (0.0)	7/65 (10.8)	0.076
mTICI ≥2b	22/27 (81.5)	49/65 (75.4)	0.596
Procedure-related complications	4/27 (14.8)	11/65 (16.9)	0.803

Values are presented as median (interquartile range) or n (%).
ASPECTS, Alberta Stroke Program Early CT Score; pmRS, premorbid modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; BP, blood pressure; CT, computed tomography; MRI, magnetic resonance imaging; ICA, internal carotid artery; GA, general anesthesia; CS, conscious sedation; sICH, symptomatic intracranial hemorrhage; mTICI, modified Thrombolysis In Cerebral Infarction.
*Significant differences.