Supplementary Table 4. Premorbid ASPECTS ≤5 patients stratified by successful recanalization

	mTICI ≥2b (n=71, 77.2%)	mTICI <2b (n=22, 22.8%)	*P*
Baseline characteristics			
Age (yr)	81.1±11.3	81.0±13.7	0.448
Female sex	51/71 (71.8)	15/21 (71.4)	0.971
pmRS	3 (3–4)	3 (3–4)	
NIHSS	17 (14–21)	19 (18–27)	0.014*
BP systolic (mm Hg)	140 (135–170)	150 (129–159)	0.125
BP diastolic (mm Hg)	78 (67–87)	87 (80–100)	0.002*
Heart rate (bpm)	85 (70–98)	87 (70–109)	0.393
Vascular risk factors			
Arterial hypertension	48/70 (68.6)	15/20 (75.0)	0.783
Diabetes	14/70 (20.0)	4/21 (19.0)	0.923
Dyslipidemia	22/70 (31.4)	6/21 (28.6)	0.804
Atrial fibrillation	33/69 (47.8)	13/21 (61.9)	0.322
Active smoking	3/62 (4.8)	2/20 (10.0)	0.591
Ship	21/71 (29.6)	7/21 (33.3)	0.790
Mothership	50/71 (69.6)	14/21 (66.7)	0.790
Workflow times (h)			
Last known normal to admission	3.6 (1.5–7.9)	3.7 (2.3–9.6)	0.703
Admission to groin puncture	1.3 (0.9–1.8)	1.6 (0.7–2.1)	0.669
Last known normal to recanalization	5.5 (3.4–10.5)	7.1 (4.5–11.1)	0.418
Stroke, imaging, and treatment characteristics			
ASPECTS	5 (3–5)	5 (4–5)	0.343
0	0/71 (0.0)	0/21 (0.0)	
1	2/71 (2.8)	0/21 (0.0)	
2	7/71 (9.9)	1/21 (4.8)	
3	9/71 (12.7)	2/21 (9.5)	
4	12/71 (16.9)	4/21 (19.0)	
5	41/71 (57.7)	14/21 (66.7)	
ASPECTS assessed on			
CT	68/71 (95.8)	19/21 (90.5)	0.320
MRI	3/71 (4.2)	2/21 (9.5)	0.320
Carotid-T	17/71 (23.9)	6/21 (28.6)	0.775
ICA (excluding carotid-T)	5/71 (7.0)	2/21 (9.5)	0.657
M1 proximal	36/71 (50.7)	6/21 (28.6)	0.086
M1 distal	10/71 (14.1)	4/21 (19.0)	0.730
M2	2/71 (2.8)	4/21 (19.0)	0.023*
Left-sided occlusion	31/70 (44.3)	9/21 (42.9)	0.908
Intravenous thrombolysis	29/71 (40.8)	5/21 (23.8)	0.202
Type of anesthesia			
GA	53/71 (74.6)	14/20 (70.0)	0.775
CS	18/71 (25.4)	6/20 (30.0)	0.775
Switch local to GA	0/71 (0.0)	0/20 (0.0)	1.000
Recanalization attempts	2 (1–3)	4 (1–6)	0.032*

https://doi.org/10.5853/jos.2025.00668

Supplementary Table 4. Continued

	mTICI ≥2b (n=71, 77.2%)	mTICI <2b (n=22, 22.8%)	*P*
Clinical and radiological outcomes			
mRS at 90 days	6 (5–6)	6 (6–6)	0.602
NIHSS at 24 h	18 (14–21)	26 (18–30)	0.578
90-day mortality	49/71 (69.0)	16/21 (76.2)	0.596
sICH	6/71 (8.5)	1/21 (4.8)	0.575
Procedure-related complications	10/71 (14.1)	5/21 (23.8)	0.320

Values are presented as mean±standard deviation, median (interquartile range), or n (%).
ASPECTS, Alberta Stroke Program Early CT Score; mTICI, modified Thrombolysis In Cerebral Infarction; pmRS, premorbid modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; BP, blood pressure; CT, computed tomography; MRI, magnetic resonance imaging; ICA, internal carotid artery; GA, general anesthesia; CS, conscious sedation; sICH, symptomatic intracranial hemorrhage.
*Significant differences.