Supplementary Table 5. Patients with ASPECTS 0–5 stratified by prestroke disability

	pmRS 3–5 (n=92; 15.2%)	pmRS ≤2 (n=513; 84.8%)	*P*
Baseline characteristics			
Age (yr)	80.5±11.9	72.2±13.6	<0.001*
Female sex	66/92 (71.7)	263/513 (51.3)	<0.001*
pmRS	3 (3–4)	0 (0–1)	<0.001*
NIHSS†	18 (15–21)	17 (14–20)	0.168
BP systolic (mm Hg)	150 (130–170)	150 (130–168)	0.775
BP diastolic (mm Hg)	80 (70–90)	80 (70–92)	0.432
Heart rate (bpm)	85 (70–99)	80 (69–92)	0.096
Vascular risk factors			
Arterial hypertension	63/90 (70.0)	376/511 (73.6)	0.520
Diabetes	18/91 (19.8)	103/511 (20.2)	0.934
Dyslipidemia	28/91 (30.8)	201/508 (39.6)	0.128
Atrial fibrillation	46/90 (51.1)	220/509 (43.2)	0.169
Active smoking	5/82 (6.1)	79/471 (16.8)	0.012*
Antiplatelet therapy‡	29/86 (33.7)	135/506 (26.7)	0.193
Oral anticoagulation§	14/86 (16.3)	60/506 (11.9)	0.298
Previous stroke	19/92 (20.7)	81/513 (15.8)	0.248
Ship	28/92 (30.4)	207/513 (40.4)	0.082
Mothership	64/92 (69.6)	306/513 (59.6)	0.082
Workflow times (h)			
Last known normal to admission	3.6 (1.6–8.0)	3.6 (1.7–8.7)	0.831
Admission to groin puncture	1.3 (0.9–1.8)	1.2 (0.9–1.7)	0.308
Groin puncture to recanalization	0.6 (0.3–1.1)	0.5 (0.3–1.0)	0.526
Last known normal to recanalization	5.8 (3.8–10.8)	5.7 (3.8–11.2)	0.972
Stroke, imaging, and treatment characteristics			
ASPECTS	5 (4–5)	5 (4–5)	0.370
0	0/92 (0.0)	0/513 (0.0)	
1	2/92 (2.2)	23/513 (4.5)	
2	8/92 (8.7)	25/513 (4.9)	
3	11/92 (12.0)	67/513 (13.1)	
4	16/92 (17.4)	128/513 (25.0)	
5	55/92 (59.8)	270/513 (52.6)	
ASPECTS assessed on			
CT	87/92 (94.6)	446/509 (87.6)	0.071
MRI	5/92 (5.4)	63/446 (12.4)	0.071
Carotid-T	23/92 (25.0)	148/513 (28.8)	0.530
ICA (excluding carotid-T)	7/92 (7.6)	28/513 (5.59	0.464
M1 proximal	42/92 (45.7)	195/513 (38.0)	0.202
M1 distal	14/92 (15.2)	71/513 (13.8)	0.745
M2	6/92 (6.5)	57/513 (11.1)	0.264
Left-sided occlusion	40/91 (44.0)	259/511 (50.7)	0.259
Intravenous thrombolysis	34/92 (37.0)	188/512 (36.7)	0.965

https://doi.org/10.5853/jos.2025.00668

Supplementary Table 5. Continued

	pmRS 3–5 (n=92)	pmRS ≤2 (n=513)	*P*
Type of anesthesia			0.147
GA	67/91 (73.6)	403/510 (79.0)	
CS	24/91 (26.4)	94/510 (18.4)	
Switch local to GA	0/91 (0.0)	13/510 (2.6)	
Recanalization attempts	2 (1–3)	2 (1–3)	0.954

Values are presented as mean±standard deviation, median (interquartile range), or n (%).
ASPECTS, Alberta Stroke Program Early CT Score; pmRS, premorbid modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; BP, blood pressure; CT, computed tomography; MRI, magnetic resonance imaging; ICA, internal carotid artery; GA, general anesthesia; CS, conscious sedation.
*Significant differences; †NIHSS at admission was assessed based on the neurological symptoms observed at the time of admission, regardless of any preexisting physical deficits from prior strokes; ‡Included medication with aspirin and/or clopidogrel; §Included medication with either apixaban, rivaroxaban, dabigatran, edoxaban, or phenprocoumon.



Supplementary Figure 1. Flowchart of patient selection. GSR-ET, German Stroke Registry–Endovascular Treatment; pmRS, premorbid modified Rankin Scale; BA, basilar artery; VA, vertebral artery; PCA, posterior cerebral artery; ACA, anterior cerebral artery; eCA, extracranial carotid artery; ET, endovascular treatment; ASPECTS, Alberta Stroke Program Early CT Score.