**Supplementary Table 3.** Distribution of hemorrhagic transformation subtypes on day 5 by glycemic trajectory group

| | Total (n=310) | Normoglycemic (Group 1, n=96) | Persistently elevated stable (Group 2, n=153) | Variable hyperglycemic (Group 3, n=61) | *P* |
|---|---|---|---|---|---|
| ECASS intracranial hemorrhage classification on day 5 | | | | | 0.009 |
| No HT | 188 (60.6) | 66 (68.8) | 83 (54.2) | 39 (63.9) | |
| Hemorrhagic infarction type 1 | 43 (13.9) | 14 (14.6) | 20 (13.1) | 9 (14.8) | |
| Hemorrhagic infarction type 2 | 41 (13.2) | 13 (13.5) | 22 (14.4) | 6 (9.8) | |
| Parenchymal hematoma type 1 | 24 (7.7) | 1 (1.0) | 21 (13.7) | 2 (3.3) | |
| Parenchymal hematoma type 2 | 14 (4.5) | 2 (2.1) | 7 (4.6) | 5 (8.2) | |

Values are presented as n (%). Data were available for 310 of the 319 patients; follow-up imaging on day 5 was not obtained in the remaining 9 patients.
ECASS, European Cooperative Acute Stroke Study; HT, hemorrhagic transformation.


https://doi.org/10.5853/jos.2026.03867